# EMMA with 4 Modules: Commissioning Similar to Full Ring

J. Scott Berg
Brookhaven National Laboratory
11 March 2010

### Outline

- Experimental tasks
  - Injection
  - Closed orbit and tunes
  - Time of flight
  - Lattice correction
  - Cavity phasing
- Advantages and Disadvantages

## Injection

- Same problem as full ring
- Can't precisely determine closed orbit
  - More on next slide
- Don't care if kicker pulse is too long
  - Allows more time to get right

#### Closed Orbit and Tunes

- Can't find truly "closed" orbit
  - Pretend cells are identical
  - Pseudo-closed orbit minimizes cell-to-cell variation
- Tunes: again, pretend cells are identical
  - NAFF on cell-by-cell data (few cells)
  - Least squares to sinusoidal

## Time of Flight

- Can't do two passes by same object
- Could use BPMs
  - Systematic offset (small) since two different BPMs
  - Still get relative variation (vs. energy)
  - Accuracy may be challenging
- Would need two (preferably identical) scopes

#### **Lattice Correction**

- Could get rough idea of tune and time of flight vs. energy
- Could correct lattice to desired configuration using this
- Correction not precisely what we want: need to check against truly closed orbit
- Likely very close, however
- Maybe correct cell-to-cell variation?

## **Cavity Phasing**

- Can't set cavity frequencies: redundant with phase when only one pass
- Can attempt cavity phasing exercise
  - Power cavities, adjust phase to minimize orbit deviation downstream
  - Won't give final settings: frequency must be set with closed ring
  - Probably waste of time, since real algorithm must do frequency and phase

### Advantage: Get Started Sooner

- Can begin getting machine in shape now
- Major injection problems identical
  - Can wait for better solution for kicker supply
  - Finding closed orbit slightly different
- Find closed orbit parameters vs. energy
  - At least good guesses
- Bring lattice closer to desired state
- Experiments not possible in ring (Shinji's talk)

## Disadvantage: Delay

- Work needed for partial ring delays work needed for full ring. Only partial overlap.
- Algorithms and procedures similar but not identical to those for full ring
  - Repeat for full ring, but will start close
- Costs in time/money
  - Additional work in construction
  - Some repetition of experimental work
  - Time better spent preparing for full ring?

## Disadvantage: Delay

- Do we demonstrate anything interesting?
  - No significant acceleration possible
  - At best almost get linear parameters vs. energy
  - Experiments that are not possible in full ring (Shinji's talk)
- Could we realistically have anything for IPAC?

### Summary

- Can begin work needed for full ring
  - Would be done imperfectly, some (maybe small) repetition needed in full ring
- There are costs associated with the delay
- Advantage of early start unclear (haven't seen Rob's talk at time of writing...)
- But partial ring may give additional experimental possibilities